



Length-Weight Relationships and Condition Factor of *Oreochromis mossambicus* (Peters, 1852) from Manchar Lake Distt. Jamshoro, Sindh, Pakistan

W. M. ACHAKZAI*, S. SADDOZAI**, W. A. BALOCH**, N. MEMON***

Department of Fresh Water Biology and Fisheries, University of Sindh, Jamshoro

Received 25th January 2013 and Revised 20th April 2013

Abstract: Oreochromis mossambicus are among the most widely distributed exotic fishes in the world and mostly used for food. The present study was undertaken to unveil information on the length-weight relationships (LWRs) and condition factor (Kn) of freshwater fish Oreochromis mossambicus (Peters, 1852) from Manchar Lake District Jamshoro, after post super flood in Sindh. Fish were studied by examining 364 specimens (Male 186 and female 78) collected from August 2011 to July 2012. These were 10-26 cm in total length (TL) and 19.8- 295 g in weight. Length-weight relationships calculated were, Log W = -1.8007+3.0556 x Log L (Combined population), Log W = -1.7802+3.0406 x Log L (male population) and Log W = -1.8062 + 3.0577x Log L (female population). The relative condition factor (Kn) for Combined population was 0.87 - 1.07 (Mean 1.0023±0.0671), while for male 0.86 - 1.09 (Mean 1.0027 ± 0.0733) and females 0.87 - 1.07 (Mean 1.0020 ±0.0631). The Length-weight relationships and condition factor indicate isometric and satisfactory growth of Oreochromis mossambicus in Manchar Lake.

Keywords: Oreochromis mossambicus , Manchar Lake, Condition factor, Pakistan

1. INTRODUCTION

Manchar Lake is the largest freshwater floodplain lake in Pakistan. It is located in the west of Indus River, in Jamshoro district, Sindh with an area from 60 Km² to as much as 350 Km².

Oreochromis mossambicus (Peters, 1852) is a true bony fish belongs to the family Cichlidae and commonly known as tilapia. It is the most common and widely distributed exotic fish in the world (Canonico et al., 2005) and mostly used as food (Gupta, and. Acosta, 2004) due to its mild white flesh attracts consumers and hence making it economically important (Courtenay, 1989). The family Cichlidae is highly diversified with a wide range of distribution spreading across Africa and most part of India and Ceylon (Balarin, 1979). This fish can live in brackish water, so imported by many countries (Mirza, 1990). O.mossambicus was first introduced in Pakistan in 1951 from Indonesia and Thailand (Frose and Pauly, 2011). In Pakistan some work on length-weight of Oreochromis (Naeem et al., 2010a, 2011a) and proximate of farmed Tilapia (O. mossombicus) is available (Naeem et al., 2011b).

The growth and condition of fish can be estimated through Length-weight relationship (LWR) and condition factor (Kn) studies (Okgerman, 2005). Length-weight relationship is helpful in conversion of growth in weight and growth in length equation to estimate the biomass from length observations and stock assessment models (Wootton, 1990; Moutopoulos and Stergiou, 2002). The aim of present study is to provide information on the Length-weight relationships (LWRs)

and Condition factor (Kn) of O. mossambicus in Manchar Lake Sindh, Pakistan.

2. MATERIALS AND METHODS

Monthly fish samples were collected from August 2011 to July 2012 from the catch of fisherman at Manchar Lake (District Jamshoro). The fish sample comprised of a total 364 specimens (186 male and 178 female) having total length (TL) 10-26 cm (males) and 10-25.6 cm (females), weighing 23.64-286.30 g (males) while females 23.85-283.67 g. Before weighing all fishes were dried using paper towels and were weighed on portable digital balance with a sensitivity of 0.1 g. Body length measurements were made to the nearest 0.1 cm using a measuring tray.

The LWRs were calculated using equation W= a L^b, where

W = total weight (g),

L = total length (cm),

a = coefficient related to body form and,

b = exponent indicating isometric growth when equal to 3.0.

The parameters a and b were estimated by linear regression on the transformed equation: Log W= log a + b log L. The smooth mean weight W, for each length group have been computed from Le Cren,s modified formula Kn = W/a L^b (Le Cren, 1951) that can be expressed as,

Kn = W/Ŵ, where,

W = observed weight and,

Ŵ = calculated weight

Corresponding author: Wazir Ali Baloch wabaloch@hotmail.com Cell +92-300-3040072

*Department of Zoology University of Balochistan, Quetta

**Department of Zoology SBK Women University, Quetta

***Department of Zoology University of Sindh, Jamshoro

Relative condition factor (Kn) for males, females and combined sexes were calculated for each 2 cm length intervals.

-1.7802, 3.0406 (males), -1.8062, 3.0577 (females) and -1.8007, 3.0556 (combined population).

3. RESULTS AND DISCUSSIONS

Length-weight relationships(LWRs)

Male length frequency distribution showed a range 10-26 cm while weight range was 19.8-295 g (Table 1). The LWRs estimated by linear regression, the equations produced are as under, **(Fig. 1)**

Log W = -1.7802+ 3.0406 Log L (Males)

In our study the value of *b* was 3.055 which is slightly different from the results of (Naeem *et al.*, 2010a) that is 2.93 whereas our results showed that almost isometric growth, where *b* = 3.05. Regression slope showed isometric growth which indicates that the small specimens have same form and condition as of large specimens (Froese, 2006; Percin and Akyol, 2009).

The female length frequency distribution showed a range 10-25.6 cm while weight range was 19.9-290 g (Table 2). The LWRs estimated by linear regression, the following equation for females was developed **(Fig. 2)**.

Log W = -1.8062+ 3.0577 Log L (Females)

The combined sex length frequency distribution showed a range 10-26 cm while weight range was 19.8-295 g. The LWRs estimated by linear regression, the equation for combined sexes **(Fig. 3)** is as follows,

Log W = -1.8007+ 3.0556 Log L (Combined)

The value of *b* in *O. mossambicus* was compared with the work of other researchers for other fish species. The results of LWRs are in general agreement with those reported by Arslan *et al.*, (2004) in *Salmo trutta* and in female *O. mossambicus* (Naeem *et al.*, 2011b). Our estimation was compared by log *a* versus *b* on fish Base (Froese and Pauly, 2011) was applied and found to be very close to those existing for *O. mossambicus*. The value of *b* was quite different from other closely related species of the same genus, *O. niloticus*, (*b* = 2.72) as reported by Naeem *et al.*, (2010 a,b; 2011a, c, d). The Variation may be due to factors like number of specimens examined, habitat, seasonal variations, health condition and differences in the observed length ranges of specimens examined (Tesch, 1971; Bagenal and Tesch, 1978; Wooten, 1998). It is also evident that pronounced seasonal fluctuations in ecological conditions result changes in condition factor due to feeding and seasonal cycles (Das, 1977).

Relative condition factor

The relative condition factor (Kn) of a size range 0.86-1.09 (SD± 0.073, mean 1.003) in males, 0.87-1.07 (SD± 0.063, mean 1.002) in females and 0.87-1.07 (SD± 0.067, mean 1.002) in combined sexes, respectively **(Table 1-3)**.

The relative condition factor showed variation in all size groups. The highest Kn values were found in smaller fishes, which is in agreement with Shafi and Quddus (1974). We conclude that the growth of *O. mossambicus* in Manchar Lake is isometric and satisfactory.

The Length-weight relationships of *O. mossambicus* are presented in Table 2. If the specific gravity and morphology of a fish remain unchanged during its life, the value of regression coefficient *b* will be exactly 3.0 in the relation $W = a L^b$ ((Ricker, 1963; Bagenal and Tesch, 1978; Wooten, 1990). In present study the values of *a* and *b* computed were,

Table 1. Descriptive statistics and estimated parameters of length-weight relationships, f (Female) m (Male) c (Combined sexes) and n (sample size).

Sex	n	Length range (cm)	Weight range (g)	a	95% CL	b	95% CL	r ²
M	186	10-26	19.8-295	-1.7802	-2.1249 -1.4356	3.0406	2.7653 3.3158	0.963
F	78	10-25.6	19.9-290	-1.8062	-2.1111 -1.5241	3.0577	2.8329 3.3021	0.963
C	364	10-26	19.8-295	-1.8007	-2.1202 -1.4812	3.0556	2.8003 3.3108	0.963

Table 2. Length –weight relationship of *Oreochromis mossambicus* in Manchar Lake, District Jamshoro.

Length Groups (cm)	Male			Female			Combined sexes		
	No.of males	Length (cm)Mean+SD	Mean weigh (g)	No.of females	Length (cm)Mean+SD	Mean weigh (g)	No.of specimens	Length (cm)Mean+SD	Mean weigh (g)
10-12	16	10.9±0.66	23.6	20	11.0±0.70	23.8	36	10.9±0.68	23.7
12-14	40	13.3±0.58	44.2	33	13.3±0.55	43.8	73	13.3±0.56	44
14-16	47	15.2±0.54	67.4	41	15.2±0.56	67.5	88	15.2±0.54	67.5
16-18	41	17.1±0.56	80.5	47	17.0±0.57	80.5	88	17.0±0.56	80.5
18-20	13	19.2±0.66	145.6	11	18.9±0.67	132.1	24	19.1±0.66	139.5
20-22	14	21.4±0.49	191.5	12	21.4±0.53	189.6	26	21.4±0.50	190.6
22-24	6	23.5±0.65	265	5	23.5±0.53	264.3	11	23.5±0.57	264.6
24-26	9	25.5±0.55	286.3	9	25.3±0.58	283.6	18	25.3±0.51	284.9
Total	186			178			364		

Table 3. Relative condition factor (Kn) for male, female and combined sexes of *Oreochromis mossambicus* at different size groups.

Length groups (cm)	Male			Female			Combined sexes		
	Observed weight(g)	Calculated weight(g)	Kn	Observed weight(g)	Calculated weight(g)	Kn	Observed weight(g)	Calculated weight(g)	Kn
10-12	23.6	23.83	0.99	23.8	24.18	0.98	23.76	24	0.99
12-14	44.2	43.64	1.01	43.8	42.7	1.02	44.05	43.16	1.02
14-16	67.4	65.07	1.03	67.5	64.6	1.04	67.53	64.85	1.04
16-18	80.5	93.02	0.86	80.5	91.78	1.04	80.5	92.49	0.87
18-20	146	132.8	1.09	132	125.9	1.04	139.5	129.81	1.07
20-22	192	185.8	1.03	190	184.7	1.02	190.7	185.68	1.03
22-24	265	246.8	1.07	264	245.8	1.07	264.7	247.07	1.07
24-26	286	313.7	0.91	284	305.3	0.92	285	308.31	0.92
Mean Kn			1.0027			1.002			1.0023



Fig. 1. Coefficient of determination of *O. mossambicus* (Male).




Fig. 2. Coefficient of determination of *O. mossambicus* (Female).




Fig. 3. Coefficient of determination of *O. mossambicus* (Combined).

REFERENCES:

- Arslan, M., A. Yildirim and S. Bektas (2004) Length-Weight Relationship of Brown Trout, *Salmo trutta* L., Inhabiting Kan Stream, Çoruh Basin, North-Eastern Turkey. *Turk. J. Fish. Aquat. Sci.* (4): 45-48.
- Bagenal, T. B. and F. W. Tesch (1978) Age and growth. In: Methods for assessment of fish production in freshwater, 3rd edn, T. Bagenal. (Ed), IBP Handbook No. 3. Blackwell Scientific Publication, Oxford, UK, 101-136.
- Balarin, A. E., (1979) Distribution patterns of cichlids in the tropical waters. Oliver, Edinburgh.
- Canonico, G. C., A. Arthington, J. K. McCrary and M.L. Thieme (2005) The effects of introduced tilapias on native biodiversity. *Aquat. Conserv.* (15): 463-483.
- Courtenay, W. R. Jr (1989). "Exotic fishes in the National Park System," in Proceedings of the 1986 conference on science in the national parks, vol. V, L.K. Thomas, Ed. Washington DC: Management of exotic species in natural communities. U.S. National Park Service and George Wright Society, 237-252.
- Das, H. P. (1977) Length – weight relationship and relative condition of grey mullet, *Mugil cephalus* L., Maha Sagar (10): 145-149.
- Froese, R., (2006) Cube law, condition factor and weight length relationship: history, meta-analysis and recommendations. *J. Appl. Ichthyol.*, (22): 241-253.
- Froese, R. P. and D., Editors (2011) Fish Base. World wide web electronic publication. <http://www.fishbase.org>. 06: 2011.
- Gupta, M.V and B.O Acosta (2004). A review of global tilapia farming practices. World Fish Center P.O. Box 500 GPO, 10670, Penang, Malaysia.
- LeCren, E.D. (1951) The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20: 201-219.
- Mirza, M.R. (1990). Freshwater fishes of Pakistan (Urdu Eds.), Urdu Science Board, 299 Upper Mall, Lahore.
- Moutopoulos, D.K. and K.I Stergiou (2002) Length-weight and length-length relationships of fish species from the Aegean Sea (Greece) *J. Appl. Ichthyol.* 18, 200-203.
- Naeem, M., A.Salam. Q. Gillani and A. Ishtiaq (2010a). Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *J. Appl. Ichthyol.* (26): 620Pp.
- Naeem, M., A.Salam. A. Ishtiaq and S. Shafique (2010b) Length- weight and condition factor relationship of farmed hybrid (*Catla catla* . x *Labeo rohita* .) from Multan, Pakistan. *Sindh Univ. Res. J. (Sci. Ser)*. 42 (2): 35-38.
- Naeem, M., A. Salam. R. Baby. A. Ishtiaq and S.A. Rasool (2011a) Study of Body Composition of Female Population of Farmed *Oreochromis mossambicus* in relation to Body Size and Condition Factor from Pakistan. International Conference on Bioscience, Biochemistry and Bioinformatics (ICBBB). 26-28, February, Singapore, 360-363.
- Naeem, M., A. Salam. M. Ashraf. R. Baby. M. Ali and A. Ishtiaq (2011b) Length-Weight Relationship of Female Population of Farmed *Oreochromis mossambicus* in Relation to Body Size and Condition Factor from Pakistan, International Conference on Bioinformatics and Biomedical Engineering, Amsterdam, The Netherlands, July 13-15, World Acad. Sci. Eng. Technol. (78): 1149-1153.

- Naeem, M., A. Salam and A. Ishtiaq (2011c) Length-weight relationships of wild and farmed *Tor putitora* from Pakistan. *J. Appl. Ichthyol.* 27 (4): 1133-1134.
- Naeem, M., A. Salam. M. Ashraf. M. Khalid and A. Ishtiaq (2011d) External morphometric study of hatchery reared mahseer (*Tor putitora*) in relation to body size and condition factor. *Afr. J. Biotechnol.* 10 (36):7071-7077.
- Okgerman, H., (2005) Seasonal variation of the length weight and condition factor of rudd (*Scardinius erythrophthalmus* L.) in Spanca Lake. *Int. J. Zool. Res.* 1, 6–10.
- Percin, F. and O. Akyol (2009) Length–weight and length–length relationships of the bluefin tuna, *Thunnus thynnus* L., in the Turkish part of the eastern Mediterranean Sea. *J. Appl. Ichthyol.* 25, 782-784.
- Ricker, W. E. (1963) Big effects from small causes: two examples from fish population dynamics. *J. Fish. Res. Board Can.* (20): 257-264.
- Shafi, M. and M. M. A. Quddus (1974) The length-weight relationship and Condition in the Carp *Catla catla* (Hamilton- Buchanan) *J. Asiatic. Soc. Bangladesh* (sc), 19 (2): 71-80.
- Tesch, F. W. (1971) Age and growth In: *Methods for assessment of fish production in freshwaters.* W.E. Ricker (Ed). *Blackwell Scientific Publications*, Oxford, pp, 99-130.
- Wootton, R. J. (1990) Ecology of teleost fishes, Chapman and Hall. London, .404Pp.
- Wootton, R. J. (1998) Ecology of teleost fishes, *Kluwer Academic Publishers, Dordrecht, The Netherlands.*